



新たに知事指定した3物質


(1) Cyputylone, N-Cyclohexylmethylone

<p>化学名 及び 構造式</p>	<p>1-(ベンゾ[d][1,3]ジオキソール-5-イル)-2-(シクロヘキシルアミノ)プロパン-1-オン及びその塩類</p>	 <p>The structure shows a benzodioxole ring system (a benzene ring fused to a 1,3-dioxole ring) at the 5-position. This is connected via a carbonyl group to a propanone chain. The alpha-carbon of the propanone chain is substituted with a methyl group and a cyclohexylamino group (-NH-Cyclohexyl).</p>
---------------------------	---	--

(2) N-Desethyl etonitazene

<p>化学名 及び 構造式</p>	<p>2-{2-[(4-エトキシフェニル)メチル]-5-ニトロ-1H-ベンゾ[d]イミダゾール-1-イル}-N-エチルエタン-1-アミン及びその塩類</p>	 <p>The structure features a 1H-benzodiazole ring system with a nitro group (-NO2) at the 5-position. At the 2-position of the benzodiazole, there is a 2-[(4-ethoxyphenyl)methyl]ethylamino group (-CH2-CH2-NH-CH2-CH2-C6H4-OEt).</p>
---------------------------	---	---

(3) 1T-AL-LAD

<p>化学名 及び 構造式</p>	<p>(8R)-6-アリル-N,N-ジエチル-1-(チオフェン-2-カルボニル)-9,10-ジデヒドロエルゴリン-8-カルボキサミド及びその塩類</p>	 <p>The structure is a complex pentacyclic alkaloid. It consists of a tetrahydroisoquinoline core fused to a benzene ring and a thiazole ring. The thiazole ring is substituted with a thiophen-2-yl group. The tetrahydroisoquinoline ring has a diethylamino group (-N(Et)2) at the 6-position and a vinyl group (-CH=CH2) at the 8-position. A carbonyl group is attached to the 1-position of the tetrahydroisoquinoline ring.</p>
---------------------------	--	--